

!"#\$%%'()(*"+, -.#"*/0102("*3.4*56**).7*3(#8)*30109% 01: 0+*9#" 8

;))())/(19&*1: 7*3(#8*7730+*90#1

<*+=>(301(&?("930+, @A0"B&4C"1D("E@F: &G*9, #"1(&HIFJK; ' &L0(3M








Biogeosciences, 15, 5991-6018, 2018

https://doi.org/10.2446

The quantitative reconstruction of past seawater salinity has yet to be achieved, and the search for a direct and independent salinity proxy is ongoing. Foraminiferal Na/Ca has great potential in this respect as Na/Ca positively correlates with salinity (e.g. Bertlich et al., 2018).



J>"

!''# \$%&' ()%*+, -./0&-)/%1+2)34/' 2+-14+.-0%&-)%&5+, 3.)/6-)/%1+ %7+*.-18)%1/, +7%&-9/1/7' &-

!:#; 2)-0./2</1=+1'>+,.'-1/1=+*&%)%,%.2+*&/%&+)%+?-@\$-+ -1-.52'2

!A# B-.'%(2)34/'2C+,3&&'1).5+B./%(B.'/2)%,'1'+D',%&42+7&%9+)<'+E'2)+B-,/7/,+E-&9+B%%.

